

## Homework 4

Due **September 24 by 11:59pm via Canvas upload**

Please show all work, writing solutions/explanations clearly, or no credit will be given. You are encouraged to work together, but everyone must turn in independent solutions; do not copy from others or from any other sources.

1. An interstellar cloud that is 50 pc thick along our line of sight absorbs 50% of the  $\lambda = 550$  nm light passing through it.
  - (a) What is the mean free path for photons of this wavelength in this cloud?
  - (b) How thick would the cloud have to be in order to absorb only 25% of the  $\lambda = 550$  nm light passing through it?
  - (c) How thick would it have to be in order to absorb 99% of the light?
  
2. The average adult human eye can be approximated as a solid sphere, with a radius of 20 mm. Human body temperature is  $98.5^\circ \text{ F} = 310 \text{ K}$ .
  - (a) In any 1 s interval, what is the total energy in your eye, due to the photons produced by your eyeball itself? Express your answer in Joules.
  - (b) Staring at a normal 100 W lightbulb that is 1 meter away results in  $8 \times 10^{-15}$  J of energy inside your eye in any 1 s interval. Which contributes more energy to the interior of your eye: the lightbulb or your eye itself?
  - (c) Why does it get dark when you close your eyes?

(HINT: Think about how the energies of the photons from the different sources are distributed.)

**CONTINUED ON NEXT PAGE**

3. In the spectrum of a star, you measure the second line of the Balmer absorption series, also called H- $\beta$ , to be at a wavelength of  $\lambda = 4862.3 \text{ \AA}$ .
- Calculate the rest wavelength of this line, in  $\text{\AA}$ .
  - Is the star moving towards you or away from you?
  - How quickly is it moving towards you or away from you?
  - Using only this information, and the fact that the star has a parallax angle of  $\pi'' = 0.1''$ , can you deduce anything about the star's sideways, or tangential, motion?
4. Assuming stars are spherical and emit radiation as perfect blackbodies:
- For a star with a surface temperature  $T = 7000 \text{ K}$ , at what wavelength are most of its photons being emitted? What visible color does this correspond to?
  - If the star has a parallax angle  $\pi'' = 0.05''$  and a measured total flux (integrated over all wavelengths, also called the bolometric flux) of  $3 \times 10^{-9} \text{ J m}^{-2} \text{ s}^{-1}$ , estimate the radius of the star relative to the Sun's radius (i.e., in solar units  $R_{\odot}$ ).
  - Is this star visible to the naked eye at a dark site? Assume all the star's flux is emitted at its peak blackbody wavelength and that the typical eye transmits  $\sim 8\%$  of the light incident on the cornea, registers  $10\%$  of the photons striking the rods and cones, the dark adjusted pupil is  $\sim 6 \text{ mm}$  wide, and that the effective integration time of the eye is about  $0.2 \text{ s}$ . (HINT: consider what signal-to-noise is needed for the star to be perceived; neglect the impact of background photons on your eye's sensitivity to the source.)
  - How many times fainter would another star be to appear to have the same apparent brightness as the above star when seen through a high-end telescope with a mirror diameter of 10 inches? (E.g.,  $10\times$  fainter corresponds to a flux of  $3 \times 10^{-10} \text{ J m}^{-2} \text{ s}^{-1}$ )