

## Homework 7

Due **October 30 by 11:59pm via Canvas upload**

Please show all work, writing solutions/explanations clearly, or no credit will be given. You are encouraged to work together, but everyone must turn in independent solutions; do not copy from others or from any other sources.

1. For stars without a distance measurement, a distance can be estimated from their spectral type and luminosity class, a method called “spectroscopic parallax.” These estimates are more accurate for some types than others; for example, luminosity class III stars on the Horizontal Branch (so named because they are arrayed horizontally on certain color-magnitude diagrams) have absolute magnitudes  $M_V = 0.5 \pm 0.2$ . Using slide #15 from the Tuesday Week 10 lectures, estimate the distance (including error bars) to the globular cluster M3.
2. In the Sun’s core, the opacity  $\kappa \approx 0.12 \text{ m}^2 \text{ kg}^{-1}$ , and the mass density  $\rho \approx 1.5 \times 10^5 \text{ kg m}^{-3}$ . In the stellar photosphere, the opacity drops to  $0.03 \text{ m}^2 \text{ kg}^{-1}$ , and the mass density to  $1.5 \times 10^{-4} \text{ kg m}^{-3}$ .
  - (a) What is the mean free path for photons in the Sun’s core?
  - (b) What is the mean free path for photons in the Sun’s photosphere?
  - (c) Photons escape the Sun via a “random walk” process, which means that every time a photon interacts with a particle, it gets scattered into a random direction. The result of this process is that, if it were to take the photon  $N$  direct steps (or scatterings, but with no change of direction) to reach the surface of the Sun from the center, it will take  $N^2$  steps for the photon to reach the surface via a random walk. Estimate how long it takes for a fusion-created photon to escape the Sun.

**CONTINUED ON NEXT PAGE**

3. At the start of the horizontal branch phase of a  $1 M_{\odot}$  star's lifetime, about 10% of the original stellar mass is in the form of helium nuclei ( $m_{\text{He}} = 6.647 \times 10^{-27}$  kg). While on the horizontal branch, the star powers itself via the "triple- $\alpha$  process," in which three helium nuclei are converted in one carbon nucleus ( $m_{\text{C}} = 1.993 \times 10^{-26}$  kg).
- How much energy is released per triple- $\alpha$  reaction?
  - What is the total energy released by fusing this amount of helium into carbon via the triple- $\alpha$  process?
  - While on the horizontal branch, the star's luminosity is  $L = 100L_{\odot}$ . If all of this luminosity is provided by the triple- $\alpha$  process in the stellar core, how long will the horizontal branch phase last?
4. Complete Step 1 in the Timeline for your Communicating Science Project. If working in a group, all members should describe their concept and possible approaches and list the other members of the group.